

## 6th foundation + Task

①

Given  $m = 1200 \text{ kg}$  ; velocity  $u = 30 \text{ m/s}$

$\therefore$  momentum  $p = m \times \text{velocity}$

$$\Rightarrow p = 1200 \times 30 = 36000 \text{ kg m/s} \rightarrow \text{D}$$

②

Given masses  $m_1 = 5 \text{ kg}$  ;  $m_2 = 25 \text{ kg}$ .

when  $p = \text{momentum}$  is constant

$$m \propto \frac{1}{v} \quad [ p = mv ]$$

$$\Rightarrow \frac{v_1}{v_2} = \frac{m_2}{m_1}$$

$$\Rightarrow \frac{v_1}{v_2} = \frac{25}{5} = \frac{5}{1} \rightarrow \text{B}$$



Let  $m_1 = \text{mass of 1st bicycle} = 10 \text{ kg}$

$$u_1 = 5 \text{ m/s}$$

$m_2 = \text{mass of 2nd bicycle} = 15 \text{ kg}$

$$u_2 = 3 \text{ m/s}$$

The momentum after collision = momentum before collision

$$= m_1 u_1 + m_2 u_2$$

$$= 10 \times 5 + 15 \times 3$$

$$= 50 + 45$$

$$= 95 \text{ kg m/s} \rightarrow A$$

(4)

Given

momentum

$$p = 18 \text{ Ns}$$

velocity

$$v = 3 \text{ m/s}$$

$$\text{mass } m = \frac{p}{v} = \frac{18}{3} = 6 \text{ kg} \rightarrow D$$

(5)

Given

$$m_{\text{ball}} = 0.4 \text{ kg}$$

$$v_{\text{ball}} = 12 \text{ m/s}$$

$$\text{Momentum} = p = m_{\text{ball}} v_{\text{ball}}$$

$$= 0.4 \times 12$$

$$= 4.8 \text{ kg m/s} \rightarrow B$$

(6)

Given

$$m = 8 \text{ kg}$$

$$\text{momentum } p = 24 \text{ Ns}$$

We know

momentum

$$p = m v$$

$$\Rightarrow 24 = 8 \times v$$

$$\Rightarrow v = \frac{24}{8} = 3 \text{ m/s}$$





⑦

Given  $m = 3 \text{ kg}$ ; Velocity =  $3\hat{i} + 4\hat{j} \text{ m/s}$ .

$$\begin{aligned} \text{momentum of the body } p &= m v \\ &= 3[3\hat{i} + 4\hat{j}] \end{aligned}$$

$$|\vec{p}| = 3\sqrt{3^2 + 4^2} = 3\sqrt{9+16} = 3\sqrt{25} = 15 \text{ m/s} \rightarrow p$$

⑧

Given mass  $m = 250 \text{ gm} = 250 \times 10^{-3} \text{ kg}$ .

Initial speed  $u = 2 \text{ m/s}$ .

Rebounding speed  $v = -2 \text{ m/s}$

$$\begin{aligned} \therefore \text{change in momentum } dp &= m(v-u) \\ &= 250 \times 10^{-3}(-2-2) \\ &= 250 \times 10^{-3} \times (-4) \\ &= -1 \text{ kg m/s} \end{aligned}$$

Advanced

①

We know momentum  $p = m v$ .

For  $p = \text{same}$   $m \propto \frac{1}{v}$

So large mass, small velocity (or) small mass, large

velocity results same momentum.

②

We know momentum  $p = m \times v$  velocity

So both mass & velocity affects momentum.



3

we know momentum = mass  $\times$  velocity.

when velocity is same

$$\text{momentum} \propto \frac{1}{\text{velocity}} \times \text{mass}$$

Since mass of truck is more than bicycle

so momentum of bicycle is more than truck.

A is false R is true  $\rightarrow$  D.

4

we know momentum = mass  $\times$  velocity.

$\therefore$  if velocity = 0 i.e. the body is at rest

then momentum is zero.  $\rightarrow$  A

6, 7, 8

Given  $m_1 = 5 \text{ kg}$ ;  $m_2 = 4 \text{ kg}$

$u_1 = -5 \text{ m/s}$ ;  $u_2 = 8 \text{ m/s}$  [Before collision]

$v_1 = 5 \text{ m/s}$ ;  $v_2 = 3 \text{ m/s}$  [After collision]

Momentum before collision =  $m_1 u_1 + m_2 u_2$

$$\Rightarrow 5 \times 5 + 4 \times 8 = 25 + 32 = 57$$

Momentum of 5 kg body after collision  $P_1 = m_1 v_1 = 5 \times 5$

$$= 25 \text{ m/s}$$

Total momentum after collision

$$P = m_1 v_1 + m_2 v_2$$

$$= 5 \times 5 + 4 \times 3 = 25 + 12 = 37 \text{ m/s}$$





## Advanced Task

9

Given  $m_{\text{car}} = 1200 \text{ kg}$  ;  $v_{\text{car}} = 25 \text{ m/s}$ .

$\therefore$  momentum of car  $= m_{\text{car}} v_{\text{car}}$   
 $= 1200 \times 25$   
 $= 30000 \text{ kg m/s}$

10

Given momentum  $= 5 \text{ kg m/s}$ .

velocity  $= 10 \text{ m/s}$ .

We know  $m = \frac{p}{v} = \frac{5}{10} = 0.5 \text{ kg}$

Task

CG's

1

According to Newton's II<sup>nd</sup> law

$$F = \frac{dp}{dt}$$

$$\Rightarrow dp = F \times dt$$

$\therefore$  Area under  $F-t$  gives change in momentum.

2

In case of vertically projected body the velocity of the body decreases gradually while it is moving up and zero at maximum height. This is due to the body moves against gravity.

Since  $p \propto v$ , momentum ( $p$ ) also decreases.





3

A shell at rest explodes, the two pieces fly in opposite direction with same momentum, due to law of conservation of linear momentum.

Let  $m$  be mass of shell, and is at rest initially

$$\text{So } u = 0$$

$$\therefore P_{\text{shell}} = M u = 0$$

After explosion  $m_1, m_2$  and  $v_1, v_2$  are masses and velocities of two pieces.

$$\therefore P_{\text{shell}} = P_{\text{pieces}}$$

$$\Rightarrow 0 = m_1 v_1 + m_2 v_2$$

$$\Rightarrow m_1 v_1 = -m_2 v_2$$

$$\Rightarrow P_{1 \text{ piece}} = -P_{2 \text{ piece}}$$

4

when no force is acting  $\therefore p = \text{constant}$

From Newton's 2nd law  $F = \frac{dp}{dt}$

when  $p = \text{constant} \therefore F = 0$

5

In case of freely falling body, velocity of the body increases gradually because the body moves in the direction of gravity

Since momentum = mass  $\times$  velocity

As velocity increases, momentum also increases



(8)

when body is at rest velocity = 0

we know that momentum = mass  $\times$  velocity

$$\therefore \text{momentum} = 0$$

(10)

we know that

$$\text{momentum } (p) = m \times \text{velocity}$$

$$\text{when } p = \text{constant} \Rightarrow m \propto \frac{1}{\text{velocity}}$$

lighter body will have more velocity.

See main level

(1)

Given mass  $m = 6 \text{ kg}$ ; velocity  $v = 2 \text{ m/s}$

$$\therefore \text{momentum } (p) = m \times v = 6 \times 2 = 12 \text{ kg m/s} \rightarrow 0$$

(2)

initial velocity  $u = -2\hat{i} - \hat{j} \text{ m/s}$

Final velocity  $v = 4\hat{i} + 7\hat{j} \text{ m/s}$

mass  $m = 2 \text{ kg}$

$$\therefore \text{change in momentum} = m(v - u)$$

$$= 2(4\hat{i} + 7\hat{j} - (-2\hat{i} - \hat{j}))$$

$$= 2(4\hat{i} + 7\hat{j} + 2\hat{i} + \hat{j})$$

$$d\vec{p} = 2(6\hat{i} + 8\hat{j})$$

magnitude of change in momentum  $|d\vec{p}| = 2\sqrt{6^2 + 8^2}$

$$= 2\sqrt{36 + 64}$$

$$= 2 \times 10 = 20 \text{ kg m/s}$$

$\rightarrow C$





3

Given mass  $m = 3M$

Initial velocity  $u = 1.5V$ ; Final velocity  $v = -1.5V$

$\therefore$  change in momentum  $dp = m(v - u)$

$$= 3M(-1.5V - 1.5V)$$

$$= 3M(-3V)$$

$$= -9MV \rightarrow B$$

-ve sign shows  $dp$  is away from wall.

4

Given mass  $m = 0.2 \text{ kg}$ ; velocity  $v = 15 \text{ m/s}$

$\therefore$  momentum of the ball  $= m \times v$

$$= 0.2 \times 15$$

$$= 3 \text{ kg m/s} \rightarrow B$$

5

Given mass  $m = 11 \text{ kg}$ ; momentum  $= 44 \text{ N s}$

We know momentum  $(p) = m \times v$

$$\Rightarrow 44 = 11v$$

$$\Rightarrow v = 4 \text{ m/s} \rightarrow A$$

6

Given  $m = 5000 \text{ kg}$ ; velocity  $v = 20 \text{ m/s}$

$\therefore$  momentum  $(p) = m \times v$

$$= 5000 \times 20$$

$$= 1,00,000 \text{ kg m/s} \rightarrow D$$



⑦

Given  $m = 500 \text{ gm} = 500 \times 10^{-3} \text{ kg} = 0.5$

Initial speed  $|u| = 4 \text{ m/s}$

Final speed  $v = -4 \text{ m/s}$

$\therefore$  change in momentum  $= m(v - u)$


$$\Rightarrow 0.5(-4 - 4)$$

$$\Rightarrow 0.5(-8)$$

$$\Rightarrow 4 \text{ kg m/s}$$

Advanced

①



Here  $v \sin \theta$  are equal and opposite so they cancel with each other.

Initial velocity  $= v \cos \theta$

Final velocity  $= v \cos \theta$

$\therefore$  change in momentum  $= m[v \cos \theta + v \cos \theta]$

$$= 2mv \cos \theta$$

Wall is at rest before and after collision so it does not its momentum is zero and does not change.

②

we know momentum  $= \text{mass} \times \text{velocity}$

As velocity doubled  $v' = 2v$

$$p \propto v \Rightarrow \frac{p'}{p} = \frac{v'}{v} \Rightarrow \frac{p'}{p} = \frac{2v}{v}$$

$\Rightarrow p' = 2p \rightarrow$  momentum doubled



Q. 7, 8, 19

Given mass  $m = 4 \text{ kg}$ ; initial velocity  $u = 0$   
Final velocity  $v = 6 \text{ m/s}$

$\therefore$  initial momentum  $P_I = m \times \text{initial velocity}$   
 $= 4 \times 0$   
 $= 0$

Final momentum  $P_F = m \times \text{Final velocity}$   
 $= 4 \times 6$   
 $= 24 \text{ kg m/s}$

change in momentum  $= P_F - P_I$   
 $= 24 - 0$   
 $= 24 \text{ kg m/s}$